

Figure 7

## Saturation Transfer Difference Spectroscopy (STD).

### Description

The STD NMR experiment (Mayer & Meyer, 1999) is another spectroscopic technique to study the interactions, in solution, between a large molecule (receptor) and a medium-small sized molecule (ligand), and, alike tr-NOESY, it is based on the Nuclear Overhauser effect and the observation and analysis of the resonances of the ligand protons.

The experiment is carried out by first registering a reference spectrum under conditions of thermal equilibrium with the irradiation frequency set at a value that is far from any ligand or protein signal (e.g. 40 ppm), i.e. the so-called off-resonance spectrum (Fig. 7, top), which is used as reference with signal intensities  $I_0$ . A second experiment is then recorded, in which the protein is selectively saturated (on-resonance spectrum; see Fig. 7, middle), giving rise to ligand signals with  $I_{\text{sat}}$  intensities. In general, the selective irradiation consists of a cascade of Gaussian-shaped pulses (low power) that saturate only a region of the spectrum that contains a few protein resonances (but not ligand signals), e.g., the aliphatic (from 0 to -1 ppm) or aromatic region (around 7 ppm), for a specific period of time (saturation time; typically from 0.5 to 5-6 seconds). The selective saturation is transferred to the whole protein via spin diffusion through the vast network of intra-molecular  $^1\text{H}$ - $^1\text{H}$  cross-relaxation pathways (intra-molecular NOE; see Fig. 7, middle), being a quite efficient processes due to the typical large molecular weight of the receptor. Also, saturation is transferred from the protein to the bound ligand via spin diffusion through inter-molecular NOEs. The dissociation of the ligand will then transfer this saturation into the bulk solution where it accumulates during the saturation time of the experiment, as a result of the much slower relaxation in the unbound than the bound state. In particular, as in fast-exchanging protein-ligand systems the enthalpic relaxation ( $R_1$ ) of fast-tumbling molecules (small) in the free state is much slower than the kinetic off-rate constant of binding ( $k_{\text{off}} \gg R_1$ ), the accumulation of ligands molecules containing some of their resonances perturbed (NOE of large molecule) results in the macroscopic detection of transferred saturation on the ligand signals in the saturated STD NMR spectrum ( $I_{\text{sat}}$ ; Fig.7, middle).




Figure 7. Scheme of the STD-NMR experiment showing the protein in surface representation and the non-exchangeable protons of the ligand as spheres. (Top) A 1D standard NMR experiments show only equilibrium intensities of the ligand in the free state ( $I_o$ ). (Middle) Upon selective saturation of some receptor signals, this is efficiently spread throughout the protein (yellow surface) by spin diffusion (intra-molecular NOEs). The fast exchange (transient binding) between the free and bound ligand states allows the transfer of magnetization (inter-molecular NOEs) from the receptor to the ligand protons in contact with the protein surface (salmon spheres, on-resonance spectrum). (Bottom) The difference

---

spectrum ( $I_0 - I_{\text{sat}}$ ) only contains the ligand signals perturbed upon binding, whose intensities reflect the proximity of each proton to the protein surface.

Furthermore, for those hydrogen atoms of the ligand establishing close contacts to the protein surface (4-5 Å) these  $I_{\text{sat}}$  values will be lower than the  $I_0$  intensities, i.e., negative inter-molecular NOE, due to the transfer of the relaxation properties of the macromolecule to the small ligand in the bound state (Fig. 7, middle). By subtracting the off-resonance from the on-resonance spectrum ( $I_0 - I_{\text{sat}}$ ) the difference or STD spectrum is obtained, which will just contain the proton signals of the ligand in close contact to the protein surface (Fig. 7, bottom), and where any signal coming from non-binding compounds is cancelled out. So, if a non-binder is present in solution its resonances will not appear in the STD spectrum. The signal intensities exclusively coming from the saturation transfer can be quantified ( $I_{\text{STD}} = I_0 - I_{\text{sat}}$ ), showing the proximity (high  $I_{\text{STD}}$ ) or distance (low  $I_{\text{STD}}$ ) of the proton to the receptor surface. Also, a blank experiment must be carried out to assure the absence of direct irradiation of the ligand. A sample containing the receptor at low concentration and a large molar excess of the ligand (1:50 up to 1:1000) is usually employed in STD NMR experiments. This precludes the perturbations of absolute STD intensities due to rebinding effects (i.e., a ligand already saturated experiences another association process, without previous full relaxation), which would impede to correctly determine the group epitope mapping (Fig. 8).

The theory of NOE indicates that the magnetization transferred from receptor to ligand protons by intermolecular NOE depends on the inverse sixth power of their distances in the bound state. Thus, the shorter the protein-ligand proton-proton distance (bound state), the stronger the intensity of the corresponding STD signal. So, by normalizing all the measured STD intensities ( $I_0 - I_{\text{sat}}/I_0$ ) against the most intense signal (which is arbitrarily assigned a value of 100%), the so-called “group epitope mapping” is obtained (expressed as percentages; see Fig. 8). This represents the fingerprint of protein-ligand contacts in the bound state, such that it illustrates which chemical moieties of the ligand are key for molecular recognition in the binding site.




Figure 8. (Left) STD growth curves (absolute values) for the recognition of Man-(1,2)-Man-O(CH<sub>2</sub>)NH<sub>2</sub> by the anti-HIV-1 human antibody 2G12. (Right) Ligand group epitope mapping, or binding epitope for Man-(1,2)-Man-O(CH<sub>2</sub>)NH<sub>2</sub> compound. Note that proton H4A receives the highest saturation (thus the 100% is arbitrarily assigned) and is used as reference to calculate the STD percentages of the other protons. Residue of mannose-A makes the main contacts with the protein in the bound state.

Binding epitopes were commonly obtained for a given saturation time, assuming that the resulting ligand group epitope mapping did not depend on the chosen saturation time. However, significantly different  $R_1$  relaxation rates of the ligand protons can produce artefacts in the epitope definition (Yan et al., 2003). In particular, protons with slower longitudinal relaxation enable a more efficient accumulation of saturation in solution such that their relative STD intensities may be significantly overestimated at long saturation times, thus also overrating the proximity of those protons to the protein surface. Indeed, the structural information that the binding epitope provides can be affected by 1) differences in  $R_1$  relaxation rates of ligand protons, 2) the extent of saturation received in the first place, and 3) the kinetics of binding. Since these sources of distortions are the consequences of differences in the ability to accumulate saturation in the free state, we can cancel them by deriving STD intensities close to zero saturation time, this is, when virtually no accumulation of saturated ligand molecules is taking place. This is usually referred to as initial slopes of the build-up curves (Fig. 9). In addition, under this approximation, possible artefacts coming from intra-molecular spin diffusion (bound state) can also be minimized. To calculate the initial slopes, Mayer and James proposed fitting the experimental build-up curves to the mono-exponential function (Mayer & James, 2004).

$$\text{Eq. 11} \quad \text{STD}(t_{\text{sat}}) = \text{STD}^{\text{max}} (1 - \exp(-k_{\text{sat}} \cdot t))$$

where  $STD(t_{sat})$  is the observed STD intensity,  $STD_{max}$  is the asymptotic maximum of the build up curve,  $t_{sat}$  the saturation time, and  $k_{sat}$  the rate constant related to the relaxation properties of a given proton that measures the speed of the STD build-up.  $k_{sat}$  and  $STD_{max}$  are derived by least-squares fit, and the initial slope ( $STD_0$ ) of the curve is obtained as

$$Eq. 12 \quad STD_0 = \left. \frac{\delta STD(t_{sat})}{\delta t_{sat}} \right|_{t_{sat} \rightarrow 0} =$$


These  $STD_0$  values are then used to characterize the binding epitope independently of T1 and rebinding effects. Note that for

When  $t_{sat} = 1/k_{sat}$  then

Eq. 13

$$STD(t_{sat}) = STD^{max} (1 - \exp^{-1}) = 0.63$$

Thus, if we plot the normalized STD factor versus the saturation time ( $t_{sat}$ ),  $k_{sat}$  can be approximated as the inverse of the interpolated  $t_{sat}$  value from a STD factor of 0.63 (Fig. 9).



This approach can be used to quantify cases of multiple binding modes and can account for the contribution of such different binding modes involving either the same ligand or different binding sites. To get protein-ligand association curves from STD NMR experiments, Mayer and Meyer introduced the conversion from observed experimental intensities ( $(I_0 - I_{sat})/I_0$ ), which depend on the fraction of bound ligand  $f_{LB}$ .

$$\text{Eq. 14} \quad f_{LB} = \frac{[PL]}{[L]_T} = \frac{[PL]}{[L] + [PL]} =$$

to STD amplification factors (STD-AF) by multiplying the observed STD by the molar excess of ligand over protein (Mayer & James, 2004.).

$$\text{Eq. 15} \quad \text{STD - AF} = \text{STD} \cdot \frac{[L]_T}{[P]_T}$$

so that the STD amplification factor (STD-AF) depends on the fraction of bound protein ( $f_{PB}$ ).  
equation\_16-2.png

Image not found or type unknown

Therefore, a plot of STD-AF values at increasing ligand concentrations will give rise to the protein-ligand binding isotherm, from which the dissociation constant  $K_D$  can be derived. However, in eq. 16 the concentration of free ligand  $[L]$  is not known. This can be solved by employing a very low total concentration of protein, such that  $[P]_T \ll K_D$ , and an excess of ligand. Under these conditions  $[L]$  can be approximated to  $[L]_T$ , so that  
equation\_17-2.png

Image not found or type unknown

and thus the dissociation constant  $K_D$  can be easily determined by plotting the normalized STD-AF values at increasing total ligand concentrations, i.e, the binding isotherm (Fig. 10). This can be done graphically by interpolation as the  $K_D$  coincides with the amount of ligand  $[L]_T$  that is necessary to reach a 50% of  $f_{PB}$  ( $f_{PB} = 0.5$ )

equation\_18-2.png

Image not found or type unknown

Therefore, the weaker the interaction, the higher the ligand concentration required to saturate half of the protein molecules.

figure10.jpg

Image not found or type unknown

The determination of  $K_D$  from STD-NMR experiments is affected by different experimental parameters (Angulo et al., 2010). In particular, the saturation time ( $t_{sat}$ ) employed (Fig. 10 and 11), the STD intensity of the signal (Fig. 11) and the fraction of bound ligand are key factors that must be chosen wisely for the accurate determination of dissociation constants. The apparent  $K_D$  increases monotonically with  $t_{sat}$ , thus underestimating the protein-ligand affinity. Similarly, the higher the STD intensity of a resonance the more overestimated  $K_D$  value will be obtained. Also, if the fraction of bound ligand is modified by increasing the receptor concentration, the apparent dissociation constant will be larger.

Therefore, the precise quantitation of the dissociation constant implies the cancelation of all the effects coming from the fast ligand rebinding process. As the effects are related to the accumulation of saturated ligand in the bulk solution, the way to fix it is to obtain the  $K_D$  value in the absence of such accumulation, i.e. monitoring the STD in the vicinity of zero saturation time. As in the cases of determination of ligand epitopes, this requires the determination of the initial slopes of the build-up curves of STD-AF values with saturation time.

The method of STD-AF initial growth rates for  $K_D$  determination involves recording a STD-AF build-up curve (STD-AF versus saturation time) for each ligand concentration, then obtaining the initial slopes (STD-AF<sub>0</sub>) and representing them graphically as a function of the concentration of added ligand calculating the  $K_D$  from the obtained exponential curves.

figure11.jpg

Image not found or type unknown

Figure 11. Effect of the experimental factors (saturation time and monitored proton) on the determination of the apparent binding constant for the BSA-L-tryptophan system on a sample of  $20 \times 10^{-4}$ M of BSA

### Category

1. News