


Eq. 5

$$K_D = \frac{[P] \cdot [L]}{[PL]} = \frac{k_{off}}{k_{on}}$$

The Equilibrium Kinetic of binding study

Description

Assuming a protein receptor P with a single binding site for the ligand L, such that they interact to transiently form the complex PL, the binding equilibrium can be described by a model of two states with a temperature dependent dissociation constant K_D :



Eq. 5

$$K_D = \frac{[P] \cdot [L]}{[PL]} = \frac{k_{off}}{k_{on}}$$

where the unimolecular dissociation rate constant k_{off} is inversely related to the lifetime of the protein-ligand complex or, equivalently, the residence time of the ligand in the bound state t_{res} (Fig. 5), whereas the bimolecular rate constant k_{on} is the probability of a productive interaction between the protein and the ligand. At any given time, the fraction of bound protein f_{PB} is given by:

$$\text{Eq. 6} \quad f_{PB} = \frac{[PL]}{[P] + [L]}$$

Combining equations 5 and 6 yields

$$\text{Eq. 7} \quad f_{PB} = \frac{[L]}{[L] + K_D}$$

This is one of the fundamental equations that drives the outcome of the NMR experiment. It shows that increasing the ligand concentration $[L]$, the fractional occupation of the receptor-binding site f_{PB} will rise in a hyperbolic function of $[L]$. In the limit $[L] \ll K_D$, the fractional population of bound protein is directly proportional to the concentration of the free ligand. In the other extreme, $[L] \gg K_D$, the protein receptor is completely saturated by the ligand ($f_{PB} = 1$). An interesting situation occurs when $[L] = K_D$, that is, the receptor is half-saturated, which means that half of the receptor molecules exist in a one-to-one complex with the ligand. Ligands with weaker affinity have larger K_D and so, to saturate the receptor-binding site, the addition of more ligand molecules is required than in the case of a ligand with higher affinity. Thus, by adjusting the receptor and ligand concentrations, it is possible to “select” the maximum K_D for an optimum observable NMR signal. This tuning of detection threshold is very useful in ligand-based screening experiments (STD NMR in particular) of large libraries of compounds.

In the free state, both receptor and ligand retain their intrinsic NMR parameters (e.g. chemical shifts, relaxation rates, translational diffusion coefficients). In each other's presence, the mutual binding affinity of ligand and receptor drives an exchange process that toggles both sets of molecules between the free and complexed states. At equilibrium, the transient presence of the ligand molecule at the protein site is responsible for chemical exchange, characterized by the exchange kinetic constant k_{ex} .

$$\text{Eq. 8} \quad K_{ex} = K_{on} \cdot [P] + k_{off}$$

All the ligand-based NMR screening experiments assume that the receptor-ligand binding is in the fast exchange limit. Indeed, this is a necessary condition for ligand-based NMR techniques to be useful, as in the fast exchange limit the equations are enormously simplified, and the total effect of the protein-ligand complex on the NMR parameter is just the weighted sum of its separate contributions (weighted average). Thus, if Q is the NMR parameter, then

$$\text{Eq. 9} \quad \langle Q \rangle = P_B Q_B + P_F Q_F$$

where $\langle Q \rangle$ is the observed exchange-averaged parameter between its free (F) and bound (B) states,

with respective populations P_F and P_B . Observed differences between Q_F and Q_B provide measurable evidence of receptor binding and often an indicator of a “hit” in screening of a large library of compounds. The ability to detect binding with adequate sensitivity depends critically on the bound state contributions ($P_B Q_B$) being significantly larger than those from the free state ($P_F Q_F$). Since typical screening conditions where $[L] \gg [P]$ are used automatically make $P_B \ll P_F$, it is therefore necessary that Q_B is amplified in the bound state (i.e. $Q_B \gg Q_F$). On the other hand, if k_{on} is well approximated by a diffusion-limited value (10^7 - 10^9 M⁻¹s⁻¹), then the slowest k_{ex} constants lie within the $10^3 < k_{ex} < 10^5$ s⁻¹ range for weak-affinity ligands (K_D in the μ M range). Since ligand-based NMR screening methods are primarily ¹H based, k_{ex} would consequently exceed most differences in intrinsic ¹H relaxation rates and rotating frame precession frequencies, thus supporting the validity of the fast exchange assumption.

Category

1. News